


NemaTrident®L



For biological control of leatherjackets (*Tipula spp*)

NemaTrident®L (Spring) contains a naturally occurring beneficial entomopathogenic nematode, *Steinernema feltiae* (15%), inert ingredients (65%) and water (20%).

Each Pack holds, 50, 250 and 500 million of *S. feltiae* infective juveniles. The nematodes are formulated in a wettable powder, enclosed in a Non-Water-Soluble pack.

NemaTrident®L (Spring) is safe, sustainable and can provide 70-80% efficacy for effective control of leatherjacket larvae at temperatures as low as 8°C if applied with Nemaspreader® - making it the most effective and versatile product currently available for combating the crane fly larvae commonly known as Leatherjackets.

Our unique Tri-Component System ensures maximum dispersal of infective juvenile nematodes within the soil which then go in search of larvae to penetrate and kill.

Directions For Use

Application rates

Application Method	Application Volume	Application	Dose/ha	Pack size 50 million	Pack size 250 million	Pack size 500 million	Number of Applications
Turfgrass	1L/m ²	Curative	2.5 billion	100m ²	500m ²	1000m ²	2-3 at weekly intervals

Application timing

Leatherjackets (Spring application)

NemaTrident®L (Spring) should be applied from March to May when older Leatherjacket larvae (L3/L4) are feeding. For additional control please see our other product NemaTrident®L (Autumn) containing *Steinernema carpocapsae*, which can also be used to control newly hatched larvae (L1/12) from late September to October .

For best results

- Use NemaTrident®L with Nemaspreader®, a wetting agent (1L/100 L of water), which ensures the effective distribution and survival of beneficial nematodes into the soil profile.
- Apply NemaTrident®L when larvae are present and close to soil surface.
- Nematodes are susceptible to ultra-violet light and desiccation. Apply during early morning or late evening during low light levels. Do not apply in bright sunlight.
- The soil surface should be moist at the time of application and if possible irrigation before application will be beneficial.
- Apply nematodes at a time when soil temperature is expected to be over 8°C for at least 3 - 6 hours per day.
- The entire pack contents must be used once opened.
- Remove all filters from nozzles on spraying equipment to avoid blockages.

continued over...

Application directions

- NemaTrident®L can be applied using conventional spraying equipment or irrigation systems.
- Where available soil injection equipment can be used to accurately deliver the nematode solution to precise depths where larvae habitat.
- Only use nozzles with at minimum diameter of 0.8mm. Pressure should not exceed 5 bar.
- NemaTrident®L should only be applied to moist soil.
- Nematodes should be applied with at least 500-1000 litres of water per hectare.
- Combining nematodes with Nemaspreader® will help the nematodes to disperse and move through the thatch layer and effectively penetrate the soil to search for and kill the leatherjacket larvae feeding on the root systems.

Ground spraying

- Clean all equipment with clean water before using NemaTrident®L.
- Remove all fine filters in the tubing and at the nozzles.
- Use required nozzles with at least 0.8mm diameter.
- Mix the nematode into 10 litres of water.
- Rinse the pack since nematodes often cling to the packaging.
- Stir the stock solution for 3-5 min until all lumps are dissolved.
- Add ¾ of the required amount of water to the 400-500 litres tank along with the required amount of Nemaspreader®.
- With the agitator running, add the required amount of NemaTrident®L through the sieve.
- Add the remaining water and begin application of the nematode maintaining agitation to avoid settlement of the nematodes until all the mixture has been applied.
- Irrigate with 1-2 liters of water per square meter after application to wash the nematodes off the grass or soil.
- As soon as NemaTrident®L is mixed with water and Nemaspreader® the entire spray solution should be used in a single application. Do not store for longer than 4 hours in tank.

Compatibility

NemaTrident®L is compatible with most chemical fertilizers and pesticides. Only a few insecticides have a negative impact on NemaTrident®L. In these cases, apply NemaTrident®L 3 days earlier or 2 weeks later. For detailed information on compatibility please visit our website: www.bionema.com.

Storage

The product may be stored for 8-10 weeks at 5 to 10°C (see expiring date on the package).

Precaution

- Keep away from children.
- Wash hands and face after handling with clean water.
- Do not freeze.

Safety

NemaTrident®L contains naturally occurring beneficial parasitic nematode and the formulation which are safe to users, consumers and the environment. There are no restrictions or safety instructions for the use of NemaTrident®L.

Limitation of liability

Efficacy of nematodes can be influenced by many factors such as storage, method of application, water volume, soil or substrate moisture and pest density. Bionema Limited cannot be held responsible for any risks. Users should follow the product instruction and contact us for help. Liability is disclaimed for insufficient quality or possible damage arising from circumstances outside of our jurisdiction.

Contact

For further assistance and expert guidance please contact us below.



Founded on research – focused on nature

Swansea University, Singleton Park, Swansea, SA2 8PP

Email info@bionema.com Tel +44 (0)1792 606 916

www.bionema.com